

- matrices involving genotype co-variates with applications in genomic prediction. *Front. Genet.* 14, 1220408. <https://doi.org/10.3389/fgene.2023.1220408>.
- Mula, W., Kurz, N. and Lemire, D. 2018. Faster population counts using AVX2 instructions. *Comput. J.*, 6, 111-120. <https://doi.org/10.1093/comjnl/bxx046>.
- Pook, T. 2020. MoBPS: Modular Breeding Program Simulator. R package version 1.4.15. <https://github.com/tpook92/mobps>.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J. and Sham, P.C. 2007. A toolset for whole-genome association and population-based linkage analysis. *Am. J. Hum. Genet.* 81, 559–575. <https://doi.org/10.1086/519795>.
- Schlather, M. 2020. Efficient calculation of the genomic relationship matrix. *BioRxiv*, <https://doi.org/10.1101/2020.01.12.903146>.
- Strassen, V. 1969. Gaussian elimination is not optimal. *Num. Math.* 13, 354–356. <https://doi.org/10.1007/BF02165411>.
- Vandenplas, J. Veerkamp, R.F., Calus, M.P.L., Lidauer, M.H., Strandén, I., Taskinen, M., Schrauf, M.F.S.G., and ten Napel, J. 2022. MiXBLUP 3.0 – software for large genomic evaluations in animal breeding programs. In *Proceedings of 12th World Congress on Genetics Applied to Livestock Production (WCGALP)*. Wageningen Academic Publishers, Wageningen, pp. 1498–1501. https://doi.org/10.3920/978-90-8686-940-4_358.
- VanRaden, P.M. 2008. Efficient methods to compute genomic predictions. *J. Dairy Sci.* 91, 4414–4423. <https://doi.org/10.3168/jds.2007-0980>.
- Walsh, B. and Lynch, M., 2018. *Evolution and Selection of Quantitative Traits*. Oxford University Press, Oxford. <https://doi.org/10.1093/oso/9780198830870.001.0001>.
- Yang, J., Lee, S.H., Goddard, M.E. and Visscher, P.M. 2011. GCTA: A tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* 88, 76–82. <https://doi.org/10.1016/j.ajhg.2010.11.011>.